



LINCOTRIMEX

INJECTABLE SOLUTION

The infectious respiratory diseases, common in hot countries, are caused in most of the cases by a great variety of bacterial agents, and due to this fact it is necessary to use different antibiotics.

LINCOTRIMEX is the result of a combination of different active ingredients. It has a high antibacterial activity and is effective against many bacterial agents as: *Haemophilus spp.*, *Pasteurella spp.*, *Streptococcus spp.*, *Staphylococcus spp.*, *Klebsiella spp.*, *Proteus spp.*, *E.coli*, *Salmonella spp.*, *Nocardia spp.*, *Corynebacterium spp.*, *Bordetella spp.* and *Clostridium spp.*

LINCOTRIMEX presents a double synergic activity with regard to the association sulfamethoxyipyridazine + trimethoprim and to the association lincomycin + spectinomycin. This double synergy ensures a higher activity than the corresponding to the addition of the activities of each drug separately.

Taking into account the bacterial agents against which each antibiotic on its own is efficient, allows us to confirm the recommendation of this speciality to treat infectious respiratory diseases.

The association with mucolytics, expectorants, antihistaminics, bronchodilators and corticosteroids, allows to reduce the characteristic symptoms of the respiratory diseases, and helps the animal to recover.

COMPOSITION

Sulfamethoxyipyridazine sodium 10 g; Trimethoprim 2 g; Lincomycin (HCl) 2,7 g; Spectinomycin (sulphate) 5,4 g; Bromhexine 230 mg; Guaicol glyceryl ether 5 g; Eucalyptol 1 g; Chlorpheniramine maleate 300 mg; Dexamethasone sodium phosphate 20 mg; Ephedrine HCl 500 mg; Excipient to 100 ml.

INDICATIONS AND TARGET SPECIES


Bovine¹, swine and ovine¹. General respiratory diseases as pneumonia, bronchopneumonia, bronchitis, etc., either if they are complicated or not with septicaemia.

¹ Animals whose milk is not destined for human consumption.

DOSAGE AND ADMINISTRATION ROUTE

By intramuscular route. In general, 2 ml per each 15 Kg b.w., twice a day.

LINCOTRIMEX EFFECTIVENESS



WITHDRAWAL PERIOD

MEAT: 10 days
MILK: Do not use

PRESENTATION

Vials of 100 ml and 250 ml.



SUPER'S DIANA, S.L.

Ctra. C-17, km 17

08150 Parets del Vallès (BARCELONA) SPAIN

Tel. +34 93 8437536 / Fax. +34 93 8436545

E-mail: export@supersdiana.com

www.supersdiana.com

03-2013 01