



DEXIANA RETARD

INJECTABLE SUSPENSION

DEXAMETHASONE is a synthetic derivative of the glucocorticoid hydrocortisone which has been used for many years in veterinary medicine. It is used in the treatment of metabolic diseases (e.g. ketosis) in ruminants, inflammatory diseases in a number of animal species and it also shows an antiallergic effect.

The hydrosoluble salts are rapidly absorbed from the inoculation site, obtaining a maximum serum concentration within few minutes after subcutaneous or intramuscular administration.

PREDNISOLONE is a synthetic glucocorticosteroid. Its pharmacological activity lasts longer than that of hydrocortisone but less than that of the longer-acting glucocorticoids such as dexamethasone. The major therapeutic uses of prednisolone are based on the anti-inflammatory and immunosuppressive activities of glucocorticoids.

COMPOSITION

- ◆ Dexamethasone isonicotinate.....100 mg
- ◆ Prednisolone.....300 mg
- ◆ Excipient to 100 ml



INDICATIONS

- **ANTI-INFLAMMATORY:** Arthritis, edemas.
- **ANTISTRESS:** Shipping fever.
- **ANTIALLERGIC:** Urticaria, eczema.

DOSAGE AND ADMINISTRATION ROUTE

By deep intramuscular route:

- **CATTLE, HORSES:** 10 ml Dexiana Retard / animal, in single dose.
- **CALVES. OVINE:** 3-5 ml Dexiana Retard / animal. in single

WITHDRAWAL PERIOD

MEAT: 8 days
MILK: 1 day

PRESENTATION

50 ml and 100 ml vials



SPECIAL PRECAUTIONS

Do not store at low temperatures.



SUPER'S DIANA, S.L.

Ctra. C-17, km 17

08150 Parets del Vallès (BARCELONA) SPAIN

Tel. +34 93 8437536 / Fax. +34 93 8436545

E-mail: export@supersdiana.com

www.supersdiana.com

03-2013/01