



# OXITOCINA DIANA

## SOLUTION FOR INJECTION

**OXITOCINA DIANA** induces contraction of smooth muscle, most importantly of the myoepithelial cells of the mammary gland, which results in milk ejection. It also results in uterine smooth muscle contraction, an effect that increases during pregnancy. It is used in the management of milk letdown and uterine contraction (induction of parturition and treatment of pyometra).

### COMPOSITION

Oxytocin (synthetic)..... 1000 IU  
Excipient to .....100 ml

### INDICATIONS

- Induction to parturition.
- Uterine inertia or atony.
- Swineagalaxia.
- Expulsion of afterbirth and residues after parturition.
- Milk letdown after parturition.
- Uterus involution after a caesarean section and alleviation of haemorrhages.
- Chronic pyometritis and endometritis to induce the exudate expulsion.
- As adjunctive therapy to antibiotics in the treatment of acute and chronic mastitis, to produce residues expulsion and to facilitate drainage.



### DOSAGE AND ADMINISTRATION ROUTE

- **OBSTETRICS** (i.v., i.m. and s.c. route):
  - **Cows:** 7.5-10 ml of OXITOCINA DIANA.
  - **Mares:** 7.5-15 ml of OXITOCINA DIANA.
  - **Sows:** 3.5-5 ml of OXITOCINA DIANA.
- **MILK EJECTION** (preferably i.v. route):
  - **Cows and Mares:** 1-2 ml of OXITOCINA DIANA.
  - **Sows:** 0.5-2 ml of OXITOCINA DIANA.

The administration can be repeated every 30 minutes, if the veterinarian considers it is necessary.



### PRESENTATION

Vials of 10 ml, 100 ml and 250 ml

### WITHDRAWAL PERIOD

MEAT: 3 days  
MILK: 24 hours



**SUPER'S DIANA, S.L.**

Ctra. C-17, km 17

08150 PARETS DEL VALLÈS (BARCELONA) SPAIN

Tel. +34 93 8437536 / Fax. +34 93 8436545

E-mail: [export@supersdiana.com](mailto:export@supersdiana.com)

[www.supersdiana.com](http://www.supersdiana.com)