



LINCOSAN

MEDICATED PREMIX POWDER

COMPOSITION

Lincomycin (HCl)..... 44 g
 Excipient to..... 1 kg

PROPERTIES OF LINCOMYCIN

- ❖ BROAD SPECTRUM OF ACTION.
- ❖ AT HIGH DOSES IT ACTS AS BACTERICIDE.
- ❖ IT IS ABSORBED VERY QUICKLY ACHIEVING PEAK SERUM LEVELS AT 2-4 HOURS FROM ADMINISTRATION.
- ❖ MINIMUM PROTEIN BINDING ⇒ GREATER ANTIBACTERIAL ACTIVITY.
- ❖ IT PRESENTS A BLOOD-BILE-INTESTINE CYCLE ⇒ LONGER TIME OF ACTION.
- ❖ IT HAS EXCELLENT TISSUE DIFFUSION ⇒ GREAT CLINICAL EFFICACY.
- ❖ IT DIFFUSES WELL THROUGH INFLAMED MENINGES ⇒ IT CAN CROSS THE ENCEPHALIC BARRIER WHEN IT IS AFFECTED BY INFLAMMATORY PROCESSES.



DOSAGE

- . For the better weight gain: 50 g/Tn of feed, for 7-10 days
- . Control of necrotic enteritis & CRD: 100 g/Tn of feed, for 7-10 days
- . Treatment of necrotic enteritis & CRD: 250-500 g/Tn of feed, for 7-10 days

WITHDRAWAL PERIOD

MEAT: 6 days

PRESENTATION

Sacks of 25 kg

SPECTRUM OF ACTION

- *Streptococcus spp.* (β -haemolytic strains)
- *Streptococcus viridans*
- *Clostridium spp.*
- *Serpulina hyodysenteriae*
- *Staphylococcus aureus*
- *Erysipelothrix r.*
- *Leptospira spp.*
- *Lawsonia intracellularis*
- *Brachispira spp.*
- *Mycoplasmas*

INFECTIOUS PROCESS

TARGET SPECIES

TREATMENT WITH LINCOSAN

INDICATIONS



- ❖ DYSENTERY
- ❖ ENZOOTIC PNEUMONIA
- ❖ In general, DISEASES PRODUCED BY SENSITIVE GERMS



SUPER'S DIANA, S.L.

Ctra. C-17, km 17
 08150 Parets del Vallès (BARCELONA) SPAIN
 Tel. +34 93 8437536 / Fax. +34 93 8436545
 E-mail: export@supersdiana.com
www.supersdiana.com